

# SOLUTIONS

1. The center of the title line above says that this exam is:

a.	Version A
b.	Version B

**Matching Table: Questions 2 through 11**

*For each numbered question fill in on the corresponding line of your NCS answer sheet the circle under the letter of the correct answer.*

E	2. First Postulate of Special Relativity	A. Equals $(1-v^2/c^2)^{-1/2}$ times $(mv)$ .
C	3. Second Postulate of Special Relativity	B. Extrapolates to zero pressure ( $P = 0$ ) at $T_C = -273^\circ\text{C}$ .
G	4. Total Relativistic Energy of a free particle	C. Light in vacuum obeys Maxwell's equations of Electrodynamics, with speed $c = (1/\mu_0\epsilon_0)^{1/2}$
A	5. Relativistic Momentum	D. Work energy plus heat energy added equals increase in internal energy
I	6. Chemical Elements	E. Reaffirms Galilean Principle of Relativity.
B	7. Ideal Gas Model	F. Measures internal energy needed to convert solid into liquid
J	8. Kelvin Temperature Scale	G. Equals sum of rest energy and kinetic energy.
D	9. First Law of Thermodynamics	H. Kinetic plus potential energy of all the individual atoms and molecules.
F	10. Latent Heat of Fusion	I Always combine into compounds in integer ratios.
H	11. Internal Energy	J. Has a zero value which can never be attained in an actual system.

**Multiple Choice**

*Insert into your NCS answer sheet the letter of the single choice that best completes the statement or answers the question.*

12. The first postulate of special relativity
- says that there is always one absolute reference frame.
  - undercuts the Galilean principle of relativity.
  - states that not all of the laws of physics are the same in every inertial reference system.
  - Does not apply Maxwell's equations of electromagnetism, which predict that light can travel through a vacuum.
  - All of the statements (a) through (d) are true.
  - None of the statements (a) through (d) is true

13. Which of the following completions, (a) through (d), of the following is **not true**? The second postulate of special relativity
- requires that the speed of light be equal to the same value,  $c = 3 \times 10^8$  m/sec, in every inertial frame. T
  - requires that the speed of light be independent of the speed of the source emitting the light T
  - requires that the speed of light be dependent upon the speed of the observer receiving the light F
  - is already implied by the first postulate of special relativity if Maxwell's equations of electromagnetism are specified to be true physical laws. T
  - None of the above completions (a) through (d) yields a true statement; i.e., all are untrue. F
  - All of the completions (a) through (d) yield true statements. F
14. In his theory of special ~~theory~~ of relativity, Einstein
- abandoned the Galilean principle of relativity. F
  - abandoned Maxwell's equations for electricity and magnetism. F
  - Showed that the ether medium for Maxwell's electromagnetic waves had to be at rest with respect to the distant stars. F
  - Postulated, redundantly to Maxwell's Equations, that the speed of light in vacuum is always  $c = 3 \times 10^8$  m/sec. T
  - postulated the Principle of Equivalence F (This is part of the General Relativity, AE's)
  - All of the above statements (a) through (e) yield true statements about Einstein's Special Theory of Gravity F
  - None of the above completions, (a) through (e) yields a true statement. F
15. On which of the following observations, (a) through (e), will two observers in different inertial systems agree about the results?
- The speed of their motion relative to one another T
  - The rate at which one another's clocks run F
  - The lengths they measure along the direction of their relative travel F
  - The synchronization of their own clocks with the moving clocks of the other frame. F
  - The simultaneity of events at separate locations. F
  - The observers will agree on none of the items (a) through (e) above.
  - The observers will agree on all of items (a) through (e) above.
16. As a space ship approaches you in outer space at a speed of  $1.2 \times 10^8$  m/sec (equal to 40% of the speed of light), its rotating beacon sends out a series of light pulses. You measure the speed of ~~the~~ each pulse to be
- $2.8 \times 10^8$  m/sec
  - $2.6 \times 10^8$  m/sec
  - $3.0 \times 10^8$  m/sec by 2<sup>nd</sup> Postulate of Sp. Rel.
  - $3.4 \times 10^8$  m/sec
  - $4.2 \times 10^8$  m/sec
  - None of the above is within 5% of the actual speed.

17. As a friend passed you at a very high speed, you exploded two firecrackers simultaneously, one at each end of her skateboard. What happened from her point of view?
- The one at the front exploded second
  - The one at the back exploded second
  - The front and back firecrackers exploded simultaneously.
  - The one at the back exploded first
  - The answer depends on the speed of the skateboard.
  - None of the above answers, (a) through (e) is a correct answer to the question.

**Scenario A (for problems 17, 18, and 19):**

A rocket ship is 220 m long when measured at rest. An observer, O, who sees the rocket ship moving past at 99.99% of the speed of light measures its length by marking the location of its nose and its tail simultaneously and then measuring the distance between the two locations,

18. What is the relativistic adjustment factor,  $\gamma = (1 - v^2/c^2)^{-1/2}$ , for the situation in Scenario A, most nearly?

- 1.0
  - 2.1
  - 7.1
  - 22.4
  - 70
  - None of the above is within 10% of the correct adjustment factor.
- $v/c = 0.9999 = 1 - 0.0001 = 1 - \epsilon$ ;  $\epsilon = 10^{-4}$   
 $\gamma = \frac{1}{\sqrt{1 - v^2/c^2}} = \frac{1}{\sqrt{1 - (1 - \epsilon)^2}} = \frac{1}{\sqrt{1 - 1 + 2\epsilon - \epsilon^2}} \approx \frac{1}{\sqrt{2\epsilon}} = \frac{10^2}{1.4}$   
 $\gamma = 70.7$

19. Suppose that the rocket in Scenario A speeds up to a constant velocity for which the relativistic factor is  $\gamma = 75$ . What length would the observer, O, in Scenario A above measure for the rocket ship, most nearly?

- 293 m
  - 220 m
  - 147 m
  - 73 m
  - 10 m
  - 3 m
  - None of the above is within 10% of the correct answer.
- $L = L/\gamma = \frac{220}{75} = 2.93 \approx 3 \text{ m}$

20. An observer,  $\bar{O}$ , on the rocket ship of Scenario A, above, watches the observer, O, measure the length of the ship. Afterwards he criticizes O's measurement by saying

- that he measured O's meter stick and found that it was in fact shorter than a meter
- that O did not actually measure position of the two ends of the ship at the same time, but that instead he first fixed the location of the front of the ship and then, afterwards, that of the back.
- that when O claimed to have insured that he was locating the front and back at the same time by firing light pulses which then arrived half way between at the same time, his light pulses were not in fact fired at the same time
- None of the above objections (a) through (c) above, is true and valid. **F**
- All of the objections, (a) through (c) above, are true and valid

21. Einstein's Principle of Equivalence is supported by the fact that
- Light is observed to be deflected when passing massive objects.
  - The gravitational mass in Newton's Law of Universal Gravitation and the inertial mass in Newton's Second Law have, within the (diminishing) experimental error, the same values.
  - The inertial pseudo-force required to explain the physics in an accelerated frame is proportional to the mass,  $m$ , of the object observed.
  - No experiment has been devised which exhibits a measurable difference between a gravitational force and an acceleration of the local frame of reference.
  - In fact, none of the "facts" cited in (a) through (d) above actually supports the Principle of Equivalence.
  - In fact, all of the facts cited in (a) through (d) above actually support the Principle of Equivalence.
22. An electron is being accelerated by a constant force to nearly the speed of light. Which of the following statements, (a) through (e), is false?
- Its kinetic energy increases steadily. **T**
  - Its relativistic speed increases at a constant rate. **F** (speed becomes nearly constant near  $c$ )
  - Its speed can approach, but not exceed, the speed of light. **T**
  - Its total energy continually increases. **T** ( $E_{TOT} = \gamma mc^2$  &  $\gamma$  increases w/  $v$ )
  - The power, or work done per unit time by the force, approaches a constant **T**  $\frac{F \Delta x}{\Delta t} = Fv \rightarrow Fc$   
 $v \rightarrow c$
  - None of the above statements (a) through (e) is false. **F**
  - All of the statements (a) through (e) above are false. **F**
23. Which of the following expressions gives the total relativistic energy of a moving object?
- $E = mc^2$
  - $E = \gamma mc^2$
  - $E = (\gamma - 1)mc^2$
  - $E = (mv^2)/2$
  - None of the above.
24. The implications of the special theory of relativity
- are true only for objects moving at very high speeds.
  - have not yet been experimentally verified.
  - apply only to tiny atomic particles.
  - cannot be tested except when speeds are comparable to  $c$ .
  - are superseded by the implications of the quantum theory of matter
  - None of the above completions (a) through (e) yields a true statement.
  - All of the above completions (a) through (e) yield true statements.
25. Superman wants to travel back to his native Krypton for a visit, a distance of  $3 \times 10^{12}$  meters. (It takes light  $10^4$  seconds to travel this distance.) If Superman can hold his breath for 500 s and travel at any speed less than  $c$ , can he make it before he suffocates?
- No, and at best he always falls short by more than 10% of the trip distance.
  - No, but at best he falls short by less than 10% of the trip distance.
  - Yes, but always just barely, with less than 1% of the trip distance to spare.
  - Yes, because he can reduce the contracted distance he travels to as small a value as he likes by setting his speed closer to that of light.
  - Yes, because for him his biological clock slows down to give him more time
  - None of the above completions yields a true statement.

26. If the inertial mass,  $m_i$ , in Newton's II law and the gravitational mass  $m_G$  in Newton's law of gravitation were NOT the same for the same object, then
- the form of Newton's law of universal gravitation would not need to be modified.
  - the form of Newton's second law would not need to be modified.
  - Einstein's prediction that the path of light is bent by gravitational fields would be invalid.
  - The Principle of Equivalence would have to be abandoned
  - All of the above completions, (a) through (d), lead to true and correct statements.
  - None of the above completions, (a) through (d), leads to a true and correct statement.
27. The law of definite volume proportions for gaseous reactants (all under the same standardized temperature and pressure conditions)
- states that equal volumes of different gases always contain equal numbers of gas particles.
  - was first proposed on purely philosophical grounds
  - follows automatically from the law of definite mass proportions
  - contradicts the fact that reactant gas volumes which combine completely to form compounds always have ratios equal to the ratios of small integers..
  - None of the above completions (a) through (d) yields a true statement.
  - All of the above completions (a) through (d) yield true statements.
28. One mole of potassium sulfide,  $K_2S$ , molecules consists of 1 mole of sulfur (S) atoms ( $A=32$ ) and 2 moles of potassium (K) atoms ( $A=39$ ). If you combine 0.5 kg of sulfur with 1 kg of potassium to make potassium sulfide, how many moles of potassium sulfide can you make?
- 39
  - 32
  - 25.6
  - 15.6
  - 12.8
  - None of the above is within 10% of the correct answer.
- No Moles of S available:  $N_S = \frac{500}{32} = 15.6$  moles S atoms*  
*" " " K<sub>2</sub> diatoms available  $N_{K_2} = \frac{1000}{2 \cdot 39} = 12.8$  Moles of K<sub>2</sub> (C)*  
*Conclude one can make 12.8 moles of K<sub>2</sub>S. Then 2.8 moles of S atoms are left over.*
29. Suppose, hypothetically, that in question 27 above, both potassium and sulfur were diatomic gases, and suppose that 2 liters of potassium combines entirely with 1 liter of sulfur to form potassium sulfide, also a gas. Then, if the chemical formula for a molecule of potassium sulfide were  $K_4S_2$ , instead of  $K_2S$ , how many liters of  $K_4S_2$  would be produced, most nearly?
- $\frac{2}{3}$
  - 1
  - $\frac{2}{3}$
  - $\frac{1}{2}$
  - $\frac{1}{3}$
  - $\frac{1}{4}$
  - None of the above is within 10% of the correct answer.
- Let 1 l of S contains N S<sub>2</sub> molecules or 2N S atoms*  
*Then it can form 2N K<sub>2</sub>S molecules, but only N K<sub>4</sub>S<sub>2</sub> molecules.*  
*Then the number of K<sub>4</sub>S<sub>2</sub> molecules formed is the same as the number of S<sub>2</sub> molecules, so that 1 l of K<sub>4</sub>S<sub>2</sub> is formed*  
*.. the same volume as the original S<sub>2</sub>: (b)*
30. Two gases are kept at the same temperature. If the molecules of gas B have 1/2 the mass of those of gas A, what is the ratio of the mean squared speed,  $\langle v^2 \rangle_{AVG}$ , of the A molecules to that of the B molecules?
- 4
  - 2
  - 1
  - $\frac{1}{2}$
  - $\frac{1}{4}$
- $T_A = T_B \Rightarrow \frac{1}{2} m_A v_A^2 = \frac{1}{2} m_B v_B^2$   
 Then  $\frac{v_A^2}{v_B^2} = \frac{m_B}{m_A} = \frac{1}{2}$  (d)

- 31 One liter of gaseous element A combines with 3 liters of gaseous element B to form 1 liter of a gaseous compound  $A_pB_q$ . If the molecules of the gases, A and B, have 2 atoms each, what is the ratio, Q/P, for the molecule  $C = A_pB_q$ ?
- a. 2/3: 2 B atoms to 3 A atoms  
 b. 1/3: 1 B atom to 3 A atoms  
 c. 3/1: 3 B atoms to 1 A atom  
 d. 3/2: 3 B atoms to 2 A atoms  
 e. 6/3 = 2: 6 B atoms to 3 A atoms  
 f. None of the above Q/P ratios would produce one liter of compound, C
- $2N_A/2N_A = V_A/V_B = 1/3 = P/Q,$   
 So that  $Q/P = 3/1$ : (C) is correct.
- 32 Which of the following is NOT assumed in our model of the ideal gas? The gas
- a. particles rebound elastically when they collide with the container wall. T  
 b. particles are always individual gas atoms. F Maybe molecules of 2 or more atoms  
 c. particles are indestructible. T  
 d. particles do not interact except when they collide. T  
 e. particles have no internal structure. T  
 f. All of the above properties (a) through (e) are properties of our ideal gas. F  
 g. In fact, none of the above properties, (a) through (e), is a property of our ideal gas. F
33. If a liter of gas has a pressure of 6.0 atmosphere, what will the pressure be if the average kinetic energy of the molecules is reduced to half its original value, while the volume is tripled?
- a. 36 atm  
 b. 18 atm  
 c. 12 atm  
 d. 6 atm  
 e. 3 atm  
 f. 2 atm  
 g. 1 atm  
 h. None of the above is within 10% of the correct value.  
 i. None of the above, because a statement about the temperature is required to specify the final pressure.
- $(KE)_i \rightarrow (KE)_f = \frac{1}{2}(KE)_i \Rightarrow T_f = \frac{1}{2}T_i$   
 Also  $V_f = 3V_i$   
 $\frac{P_f V_f}{P_i V_i} = \frac{CT_f}{CT_i} \Rightarrow \frac{P_f}{6.0} \cdot \frac{3}{1} = \frac{1}{2} \quad \text{i.e. } P_f = \frac{6.0}{3.2} = 1.0 \text{ atm (g)}$
- 34 The two fixed points used to define the modern Celsius temperature scale are those of
- a. boiling water and a mixture of ice and salt.  
 b. the human body temperature and a mixture of ice and salt.  
 c. the human body temperature and freezing water.  
 d. boiling water and freezing water.  
 e. None of the above
- 35 What Celsius temperature corresponds most nearly to  $-40^\circ\text{F}$ ?
- a.  $8^\circ\text{C}$   
 b.  $0^\circ\text{C}$   
 c.  $-4.5^\circ\text{C}$   
 d.  $-72^\circ\text{C}$   
 e.  $-40^\circ\text{C}$   
 f. None of the above answers is within 10% of the correct answer
- $T_C = (T_F - 32) \cdot \frac{5}{9}$  yields  $10^\circ\text{ at } 32^\circ\text{F}$   
 $T_C = (-40 - 32) \cdot \frac{5}{9} = -40^\circ\text{C}$   
 yields  $100^\circ\text{ at } 212^\circ\text{F}$

36 Which of the following doubles with a doubling of the absolute temperature of a given ideal gas?

- a. average momentum
- b. average speed
- c. average velocity
- d. average mass

$$\langle KE \rangle \propto T_A \Rightarrow \langle KE \rangle \text{ doubles with } T_A.$$

$$\langle KE \rangle = \left\langle \frac{1}{2} m v^2 \right\rangle \propto \langle v^2 \rangle_{\text{avg.}}$$

- e. average of the square of the speed
- f. All of the quantities, (a) through (d) above, double with the absolute temperature.
- g. None of the quantities, (a) through (d) above, doubles with the absolute temperature

37 Which of the following doubles with a doubling of the Celsius temperature of an ideal gas?

- a. average momentum
- b. average speed
- c. average kinetic energy
- d. product of pressure and volume
- e. All of the quantities, (a) through (d) above, double with the Celsius temperature.

- f. None of the quantities, (a) through (d) above, doubles with the Celsius temperature

38 The pressure in a rigid container filled with gas increases when it is heated because

- a. the walls must do more work on the gas as  $T$  increases. **F**
- b. the volume of the gas increases with temperature. **F**
- c. the number of gas particles increases with temperature. **F**
- d. the walls must exert a larger impulse to turn back the particles. **T**
- e. All of the above completions (a) through (d) yield true statements. **F**
- f. None of the above completions (a) through (d) yields a true statement. **F**

39. Avogadro suggested that each liter of any two gases under identical conditions has the same

- a. mass
- b. number of atoms
- c. number of molecules
- d. density
- e. number of gas particles
- f. None of the above.

40. What happens to the volume of an ideal gas if you double its pressure and reduce its absolute temperature by one-half? The volume

- a. It quadruples
- b. It doubles
- c. It stays the same.
- d. It is cut in half
- e. It is cut to one-fourth its original value

$$\frac{V_f \cdot P_f = \frac{4}{1} T_f}{V_i \cdot P_i = \frac{1}{2} T_i} \Rightarrow \frac{V_f}{V_i} \left( \frac{2}{1} \right) = \left( \frac{1}{2} \right)$$

$$\text{or } V_f/V_i = 1/4$$

- f. None of the above is within 10% of the correct answer.

41. In radiative heat transfer, thermal energy is transported by
- physical vibrations of the intervening medium. **F**
  - the collisions of particles. **T**
  - the movement of a fluid. **T**
  - the propagation of sound waves. **T**
  - Electromagnetic fields. **T**
  - All of the mechanisms (a) through (e) above transmit thermal energy in radiative transfer. **F**
  - None of the mechanisms (a) through (e) above transmits thermal energy in radiative transfer. **F**
42. Joule's experiments with hanging weights turning paddle wheels in water
- showed that mechanical energy was converted to internal energy by viscous forces. **T**
  - showed that 4.2 joules of work are equivalent to 1 calorie of heat. **T**
  - fixed the ratio between the Joule and the (independently defined) calorie. **T**
  - led to a generalization of the law of conservation of mechanical energy. **T**
  - None of the above completions, (a) through (d) provides a true statement. **F**
  - All of the above completions, (a) through (d) provide true statements. **F**
43. Two objects are in thermal equilibrium if
- are in thermal contact and there is no net flow of thermal energy. **T**
  - they have the same internal energy per unit mass. **F**
  - they have the same internal energy. **F**
  - they have the same specific heat. **F**
  - All of the above completions, (a) through (d) provide true statements. **F**
  - None of the above completions, (a) through (d) provides a true statement. **F**
44. Which of the following completions, (a) through (d) below, leads to a false statement? The first law of thermodynamics
- guarantees that if work is done on a system in some process and no net heat is ejected, the internal energy of the system must increase. **T**
  - is a restatement of the law of conservation of energy. **T**
  - requires that heat can be completely converted into work. **F**
  - treats heat as another form of energy. **T**
  - All of the above statements (a) through (d) are true of the first law: None is false. **F**
  - None of the above statements (a) through (d) is true of the first law: All are false. **F**
45. If during some process a system has no change in internal energy, we can, because of the first law of thermodynamics, conclude that
- the system ejected no heat. **F**
  - no work was done on the system. **F**
  - the net amount of work done by the system was equal to zero. **F**
  - the system received no heat. **F**
  - that any net heat transfer was cancelled by the net work done. **T**
  - None of the above assertions can be made on the basis of this statement. **F**
  - All of the above assertions are guaranteed by the condition stated. **F**

46. Which of the following statements about a cup of water and a gallon of water at the same temperature is correct?
- They both will transfer heat energy under contact to a third object at lower temperature. **T**
  - They have the same internal energy per unit mass. **T**
  - The average molecular speed in the gallon is the same as that in the cup. **T**
  - The total internal energy in the cup of water is less than that in the gallon. **T**
  - The average kinetic energy of a molecule is the same for both the cup and the gallon. **T**
  - None of the above statements, (a) through (e) is true. **F**
  - (g)** All of the above statements, (a) through (e) are true. **T**
47. When an ideal gas was compressed and then allowed to expand, its internal energy increased by 10 J, it gave off 20 J of heat, and in expanding performed 30 J of work. How much work was done on the gas in the original compression?
- $$W_{IN} + Q_{IN} - W_{OUT} - Q_{OUT} = \Delta U \quad (1^{st} \text{ LAW})$$
- $$W_{IN} + 0 - 30 - 20 = 10 \text{ J}$$
- $$W_{IN} = 60 \text{ J} = (10 + 20 + 30) \text{ J}$$
- 10 J
  - 20 J
  - 30 J
  - (d)** 60 J
  - 110 J
  - None of the above is within 10% of the correct answer.
48. The third law of thermodynamics
- is a restatement of the law of conservation of energy. **F**
  - says that heat cannot be completely converted to mechanical energy. **F**
  - guarantees that temperature is useful for predicting heat transfer. **F**
  - says that all motion ceases at absolute zero. **F**: 3<sup>rd</sup> LAW does NOT address motion of particles.
  - (e)** says that we can never reach the absolute zero of temperature. **T**
  - None of the above completions correctly characterizes the third law. **F**
49. Why do climates near the coasts tend to be more moderate than near the middle of the continent?
- Because water has a high latent heat of vaporization.
  - Because the coasts have lower elevations, and cool air flows downhill.
  - Because water has a relatively high latent heat of fusion
  - Because it rains a lot on the coasts.
  - (e)** Because water has a high specific heat, and can therefore remove much heat from hotter air & supply much heat to cooler air with relatively small changes to its own temperature.
  - None of the above.
50. In our laboratory measurement of the specific heat of copper, a hot copper cylinder is immersed in a cup of ice cold water, and allowed to come to equilibrium. During this process heat transfer to or from the equilibrating (copper + water) system may cause the result to be erroneous. For our experimental procedure, it is clear that such heat transfer is most likely to lead to net heat's being transferred
- into the system, and therefore to an increase in the computed specific heat of Cu.
  - out of the system, and therefore to an increase in the computed specific heat of Cu.
  - into the system, and therefore to a decrease in the computed specific heat of Cu.
  - out of the system, and therefore to a decrease in the computed specific heat of Cu.
  - (e)** In fact, it is not clear whether net heat will be transferred in or out of the system, so that the effect on the computed specific heat can not be predicted reliably.

51. If it takes about 6600 cal to raise the temperature of a 700-g metal statue by 165° C, of which of the following materials is the statue most likely composed?

- a. Aluminum, with specific heat  $c = 0.215 \text{ cal/g}\cdot^\circ\text{C}$
- b. Copper, with specific heat  $c = 0.092 \text{ cal/g}\cdot^\circ\text{C}$
- c. Gold, with specific heat  $c = 0.031 \text{ cal/g}\cdot^\circ\text{C}$
- d. Silver, with specific heat  $c = 0.057 \text{ cal/g}\cdot^\circ\text{C}$
- e. None of the above is within 10% of the correct answer.

$$\begin{aligned} \text{Specific Heat} = c &= \frac{Q}{m \Delta T} \\ &= \frac{6600}{700 \cdot 165} = 0.0571 \frac{\text{cal}}{\text{g}\cdot^\circ\text{C}} \end{aligned}$$

52. In convective energy transfer, thermal energy is transported by

- a. the propagation of sound waves .
- b. the collisions of particles.
- c. electromagnetic fields.
- d. the movement of a fluid.
- e. physical vibrations of the intervening medium.
- f. All of the mechanisms (a) through (e) above transmit thermal energy in convection.
- g. None of the mechanisms (a) through (e) above transmits thermal energy in convection.

53. In one lab experiment one's touching of panels of various materials setting on the table suggested that the steel panel was cooler than the styrofoam panel, despite the fact that the thermometer measured the same temperature for both (and reasonably so, since they had plenty of time to come to temperature equilibrium with the air in the room). If one knew also that steel was a much better heat conductor than styrofoam one might reasonably infer that

- a. that our touch may be measuring not the temperature, but the rate of heat transfer.
- b. that styrofoam stays hot longer than steel.
- c. that our sense of touch is a good built-in thermometer.
- d. that just because an object is hot to touch it may still not be dangerous.
- e. All of the completions (a) through (d) above provide reasonable inferences.
- f. None of the completions (a) through (d) above provides a reasonable inference.

54. The first law of thermodynamics, which specifies that energy in the form of heat or work transferred between the outside world and a material system alters the internal energy of the system, is valid only

- a. if no work is done on the system. **F**
- b. when there is no friction. **F**
- c. when all of the forces acting are conservative. **F**
- d. if there is no heat loss or gain. **F**
- e. if the third law of thermodynamics is valid. **F**
- f. Each of the above completions (a) through (e) yield a true statements. **F**
- g. None of the above completions (a) through (e) yields a true statement. **T**

The following six problems #55 - #60 may require more computation than average, and you may wish to allocate your effort accordingly.

55. A super-train is traveling along a straight, horizontal track at a constant speed,  $V=0.97c$ . It fires a super-rocket in the forward direction with a speed,  $v'=0.95c$ . (Recall that, relativistically,  $S$  in  $O$  observes  $v = (v'+V)/(1+Vv'/c^2)$ ), when  $S'$  in frame  $O'$ , moving with  $V$  in  $O$ , observes  $v'$ .) An observer in the train station will measure the speed of the rocket, most nearly, to be

- a.  $1.92c$
- b.  $1.71c$
- c.  $1.0c$
- d.  $0.97c$
- e.  $0.96c$
- f.  $0.95c$
- g.  $0.52c$
- h.  $0.02c$
- i.  $0.00c$

$$v = \frac{(0.97 + 0.95)c}{1 + (0.97)(0.95)} = 0.9992c$$

is within 1% of  $c$ .

- j. None of the above is within 1% of the correct speed.

56. If a particle of rest mass,  $m=1$  kg, is accelerated from rest to a speed of  $v = 0.999c$ , what is its final kinetic energy (=total energy - rest energy), most nearly?

- a.  $9 \times 10^{20}$  J
- b.  $2 \times 10^{20}$  J
- c.  $9 \times 10^{18}$  J
- d.  $2 \times 10^{18}$  J
- e.  $9 \times 10^{16}$  J
- f.  $2 \times 10^{16}$  J
- g. None of the above is within 10% of the correct answer.

$$(KE) = (\gamma - 1)mc^2 = (22.36)1 \cdot (3 \times 10^8)^2 = 201 \times 10^4 \text{ J} = 2.0 \times 10^{18} \text{ J}$$

$$\text{Since } \gamma = \frac{1}{\sqrt{1 - (0.999)^2}} = \frac{1}{\sqrt{1 - (1 - 10^{-3})^2}} = \frac{1}{\sqrt{2 \cdot 10^{-3}}} = 22.36$$

57. One liter of an ideal gas is heated from 27 °C to 927 °C while the pressure increases from 1 atmosphere to 1.33 atmospheres. What is the final volume of the gas, most nearly?

- a. 0.25 liter  
 b. 0.33 liter  
 c. 0.50 liter  
 d. 1 liter  
 e. 2 liters  
 f. 3 liters  
 g. 4 liters  
 h. None of the above is within 10% of the correct answer

$$27^{\circ}\text{C} = 300\text{K}; 927^{\circ}\text{C} = 1200\text{K} \quad P_f/P_i = 3$$

$$\frac{P_f V_f}{P_i V_i} = \frac{T_f}{T_i} \Rightarrow (1.33) \frac{V_f}{1} = \frac{1200}{300} \Rightarrow V_f = \frac{4}{1.33} = 3 \text{ l}$$

58. A hypothetical balloon filled with an ideal gas has a volume of  $10^5$  liters at 30°C under one atmosphere of pressure. At what temperature, most nearly, will its volume be  $10^6$  liters under ten atmospheres of pressure?

- a. 3°C  
 b. 30°C  
 c. 300°C  
 d. 3000°C  
 e. 30,000°C  
 f. 300,000°C

$$\frac{P_f V_f}{P_i V_i} = \frac{CT_f}{CT_i} \Rightarrow \left(\frac{10}{1}\right) \left(\frac{10^6}{10^5}\right) = \frac{T_f}{303\text{K}} \Rightarrow T_f = 30,300\text{K}$$

$$= (30,300 - 273)^{\circ}\text{C}$$

$$= 30,027^{\circ}\text{C} \text{ (e)}$$

$$T_c = 30^{\circ}\text{C} = 303\text{K} = T_A$$

- g. None of the above is within 10% of the correct Celsius temperature

59. The latent heat of fusion for water is 80 cal/g, and its specific heat is 1 cal/g-°C. How much heat would it take to melt 3 g of ice at 0° C to form water at 70° C?

- a. 3 cal
- b. 70 cal
- c. 150 cal
- d. 210 cal
- e. 240 cal J
- f. 450 cal J

$$Q = 3 \cdot 80 + 3 \cdot 1 \cdot 70 = 240 + 210 = 450 \text{ cal}$$

g. None of the above is within 10 % of the correct answer.

60. If 100 g of steam at 100°C and 500 g of ice at 0°C are enclosed in an insulated container, what is the final equilibrium temperature, most nearly? (Recall that the latent heat of fusion of water is 80 cal/g, and the latent heat of vaporization is 540 cal/g, and use 1 cal/gm °C for the specific heat of water).

- a. 40° C
- b. 50° C
- c. 60° C
- d. 70° C
- e. 80° C
- f. 90° C
- g. 100° C

$$(100 \text{ g})(540) + (100 - T_f)(100)(1) - (500)80 - (500)(1)(T_f - 0^\circ) = 0$$

$$\text{(Heat out of Steam)} - \text{(Heat into Water)} = 0$$

$$(100)(540 + 100 - 400) = 600T_f$$

$$240 = 6T_f$$

$$40^\circ \text{C} = T_f$$

h. None of the above is within 10% of the correct answer.

.....End of Exam