

## Lab 7: How do charged objects in a fluid interact with each other and respond to external electric fields? Electrophoresis and Charge Screening in Fluids.

### Introduction


The biophysical motivation for this laboratory is the ubiquity of charges in living systems. Large molecules—in particular, proteins—are often phosphorylated and thus carry net charge. These charges generate electric fields, exerting attractive forces on other molecules, and playing a role in the intricate balance of forces and movements that occur in a living cell! We might ask: how far is the reach of the force field from a charge in a large protein when many ions are present in the surrounding fluid? How fast would a charged protein move in an electric field? These effects are crucial for understanding biological systems at the cellular level.

In this two-week lab sequence, we will investigate related questions in a simple model system. You will be investigating charge screening in fluids by employing the technique of electrophoresis. Your investigation will come in two parts: Part 1, an investigation of glass beads in de-ionized water (DI water, pure  $\text{H}_2\text{O}$ ); and Part 2, an investigation of glass beads in two saline solutions of different concentrations.

We have been employing solutions of glass beads in water in most of our labs. You may have noticed that some of the beads are stuck together, forming clumps (also called 'flocs' because their *aggregation* (coming together) is a form of *flocculation*), or that the beads sometimes "settle-out" of the fluid (like the *sedimentation* we observed in the tilted microscope lab last semester). Aggregation (flocculation or coagulation) and sedimentation are behaviors common to *colloidal* fluids. A colloidal system is one in which one phase of matter is finely dispersed throughout another phase of matter. Our glass beads in water are colloidal fluids because the glass beads (solids) are small particles dispersed throughout the water (fluid). In fact, when the glass beads are submerged in water, they become charged—even in DI water! The glass beads,  $\text{SiO}_2$ , have surface groups of  $\text{SiOH}$ . When immersed in water, the hydrogen nucleus breaks free (increasing the pH of the fluid due to the roaming  $\text{H}^+$ ) and leaving an  $\text{SiO}^-$  behind. Thus the surface of the glass beads becomes negatively charged—each bead carries charge on the order of femto-Coulombs (fC). (Lest you think a fC is small, **about how many extra electrons are on each bead?**) [You might then ask, if all the beads are negatively charged, then why do they stick together? Besides the electric repulsive forces between the beads, there are also attractive van der Waals forces; at close enough distances, the attraction dominates the interaction and the beads will stick together to form a floc.] If a salt (like  $\text{NaCl}$ ) is added to the fluid, it will dissolve and the free cations (or anions, for a positively charged colloid) will group around the negatively charged beads of glass, thus decreasing the 'effective charge' of the unit—this is called 'charge screening' or 'Debye screening.' The more ions that are available in the fluid, the greater the charge screening effect will be; thus the concentration of the electrolyte is important.

## Investigation

To investigate this charge screening effect, we will need to determine how charged our glass beads are in DI water (Part 1) and then compare that charge to the 'effective charge' as seen in various concentrations of saline solution (Part 2). We can investigate these charges using the technique of electrophoresis. By applying an electric field (generated by a potential difference between two electrodes) to the fluid (see picture above right), we can cause an electric force on the charged beads that will induce motion through the still fluid. A larger electric field will cause faster motion. **Before you begin Part 1**, you will need to model the situation: consider what forces act on the bead as it moves through the fluid and determine how changing the potential difference between the electrodes and measuring the resultant speed of the beads will enable you to find the charge on the beads. Carefully reflect on what assumptions you are making as you model the situation and think about the implications these assumptions have for the design of your experiment.



The experimental set-up available (see picture below right) includes:

- the microscope (with camera),
- a power source (the big box on the right with knobs and wires) to create constant voltage of your choosing,
- two wires with banana plugs on one end and micro-grabbers (for holding electrodes, see larger images in photo above) on the other,
- short segments of copper wire to use as electrodes,
- an eight-well chamber slide (fill one of these chambers halfway with solution for each investigation—when investigating a different solution, you can simply move to another chamber, or choose to empty the chamber and rinse with distilled water. At the end of the day please rinse the chamber with distilled water),
- three vials of solution: one of distilled water and two different saline solutions (LOW and HIGH), and
- other tools (rulers, paper towels, pipettes, etc.).



If you have never used a power source before, ask your TA for safety and use instructions. Here are some other details that may help you design your experiment:

- The maximum voltage the power source can produce is 18 V; you will likely need 4 V or more to see bead motion in DI water.
- The chambers each hold less than 1 mL of fluid; the vials provided should be sufficient.
- All solutions contain glass beads that are 5  $\mu\text{m}$  in diameter.

- The low concentration saline solution is 9 mg/L NaCl; the high concentration saline solution is 90 mg/L NaCl. The viscosities of these solutions are almost identical to that of pure DI water— $9.0 \times 10^{-4}$  Pa-s at 26 °C.
- You should gather your video data as soon after applying the potential difference to the electrodes as possible (think about why this might be so).
- You should be careful to leave the fluid sample on the microscope stage (above the hot bulb) for as short a time as possible before gathering your video (think about why this might be so).
- You may not need to use particle tracking (manual or automated) for these videos; **look for the hint** at the end of this lab.
- Though you may not need to track, you will need **lots** of data—collect carefully and precisely, collect sufficient sets of data, and consider ways to reduce your uncertainty.

### Part 1: DI Water

Design an experiment to determine the charging of beads in fluid. Consider 'broad stroke' details (e.g., What data are we collecting? How much data is 'enough?') as well as 'fine stroke' details (e.g., What part of the slide are we viewing? How are we collecting data? How do our assumptions affect our procedure and analysis?). Be sure to consider the qualitative aspects of the motion, as well (e.g., Is this really directed motion? Could it be random motion? Which electrode (+ or -) are the beads moving toward?). Once you have a good experimental design, gather data to determine the charging of our glass beads in DI water. **Be sure to note the TIMING used by the video capture program, VirtualDub, for EACH video taken.**

### Part 2: Saline Solutions of Low and High Concentration

Now investigate how saline solutions 'screen' the charge on the glass beads. Using your experimental design from Part 1, find the 'effective charge' of our glass beads in a LOW concentration saline solution. Examine the effect of a HIGH concentration saline solution. **Before you do either of these**, it may help to think about a model for a charged glass bead in an electrolyte solution.

**\*\*\* Be sure that each group member has a copy of all data- and word-processing documents created before leaving the lab at the end of week 1. \*\*\***

### **ImageJ Analysis Hint:**

Manual tracking or automated tracking may be 'overkill' for these videos. If you wish, the velocity of a single bead can be determined by measuring the displacement over a corresponding time interval. (What assumptions are implicit in this method? Are they justified assumptions? Under what conditions?) Select a bead and note both the initial frame at which it appears in your video and the final frame before it disappears from your video. In the initial frame, use the 'point tool' (plus shape icon) to click on the location of the bead. Use the scroll bar to advance the video

to the final frame; note that the location you clicked on with the 'point tool' remains visible. Using the 'line selection tool' (slanted line icon), draw a line segment from the location left by the 'point tool' to the location of the bead in the final frame. Note the pixel length of this line. (If you let go of the mouse too quickly to note the line's length, you can always choose 'Analyze,' 'Measure' to find the length of the line segment.) Using appropriate pixel to distance conversions and frame to time conversions, you can now state the velocity of this glass bead.

### For the Lab Report:

Beyond the expectations of a normal lab report (journal, procedure, data, analysis, conclusions, etc.), be sure to include careful statements of the models you have developed and the assumptions contained within those models. Also include qualitative descriptions of observed phenomena to accompany the quantitative analysis of each colloidal fluid. Include as much procedural detail, data, and analytical detail as you feel are necessary to prove your points and substantiate your conclusions. Discuss the ways in which the ideas explored here relate to biology and/or chemistry.

### Approximate Timing:

#### Week 1:

- Introduction ..... 10 min.
- Modeling Situation ..... 15 min.
- Designing Experiment ..... 15 min.
- Gathering Part 1 Videos ..... 20 min.
- Gathering Part 2 Videos ..... 30 min.
- Analyzing Videos in ImageJ ..... 15 min.
- Start Analyzing Data in Excel ..... 5 min.

#### Week 2:

- Introduction ..... 10 min.
- Finish Analyzing Data in Excel ..... 30 min.
- Creating Posters/Presentations ..... 15 min.
- Posters/Presentations ..... 25 min.
- Class Discussion ..... 10 min.
- Finalizing Lab Reports ..... 20 min.