

Physics 601 Homework 11---Due Friday Dec. 5

1. This problem concerns force on objects in rotating frames
 - a. Consider a particle which is constrained to the surface of a cylinder of radius R rotating about its axis with an angular velocity ω . A position on the cylinder is specified by z and θ (which by construction are body fixed.)
 - i. Suppose a particle of mass is moving with speed v along a path with fixed θ , that is up the cylinder. What is the magnitude and direction of the force (excluding the force of constraint) on the particle?
 - ii. Suppose a particle of mass is moving with speed v along a path with fixed z , that is around the cylinder. What is the magnitude and direction of the force (excluding the force of constraint) on the particle?
 - b. Consider a particle which is constrained to the surface of a sphere of radius R rotating about an axis through the center with an angular velocity ω . A position on the cylinder is specified by a polar, angle θ and an azimuthal angle ϕ (which by construction are body fixed.)
 - i. Suppose a particle of mass is moving with speed v along a path with fixed θ , that is along a line of latitude. What is the magnitude and direction of the force (excluding the force of constraint) on the particle?
 - ii. Suppose a particle of mass is moving with speed v along a path with fixed ϕ , that is along a line of longitude. What is the magnitude and direction of the force (excluding the force of constraint) on the particle? (Note this may depend on θ .)
2. Consider the application of the formal short time expansion developed in class to a simple Harmonic oscillator: $L = \frac{1}{2}m\dot{x}^2 - \frac{1}{2}\lambda^2 m\omega_0^2 x^2$ with $x = x_0 + \lambda x_1 + \lambda^2 x_2 + \dots$. Suppose the initial conditions are fixed by $x_0(0) = x(0); \dot{x}_0(0) = 0; x_1(0) = 0, \dot{x}_1(0) = \dot{x}(0)$ with $x_n(0) = 0, \dot{x}_n(0) = 0$ for $n > 1$. λ is taken to unity at the end of the problem.
 - a. Solve for $x(t)$ to 5th order.
 - b. Verify that the solution is identical to the exact solution Taylor expanded in time up to t^5 .
3. In class we formally developed the expansion for a system with one degree of freedom. In this problem I would like you to do it for 2 degrees of freedom : $L = \frac{1}{2}m(\dot{x}^2 + \dot{y}^2) - \lambda^2 V(x, y)$ with $x = x_0 + \lambda x_1 + \lambda^2 x_2 + \dots ; y = y_0 + \lambda y_1 + \lambda^2 y_2 + \dots$. Choosing sensible boundary conditions find an expression $x(t), y(t)$ up to 4th order. The result should be given in terms of the initial position, the initial velocity, and partial derivatives with respect to V evaluated at the initial position.

4. Apart from short time expansion there is a context in which one can build an expansion in which the full potential is treated perturbatively: if one is working in a regime in which potential energy difference are always much less than the kinetic energy the approximation can be justified up to comparative long times. Consider the case of a physical pendulum whose Lagrangian is given by $L = \frac{1}{2} I \dot{\theta}^2 + V_0 \cos(\theta)$ where I is the moment of inertia and V_0 is the maximum value of the potential energy (i.e. $m g L$) where L is length from the pivot point to the center of mass. The problem under consideration is this: suppose that at $t=0$ the system is at the minimum of the potential ($\theta=0$) with an initial angular velocity $\dot{\theta} = \omega_0$.

To develop the expansion insert powers of λ : $L = \frac{1}{2} I \dot{\theta}^2 + \lambda V_0 \cos(\theta)$ with $\theta = \omega_0 t + \lambda \theta_1 + \lambda^2 \theta_2 + \dots$ and impose boundary conditions $\theta_i(0) = 0$, $\dot{\theta}_i(0) = 0$ for i . These boundary conditions are designed to ensure that the boundary conditions to our problem is solved

Find explicitly the form for θ_1 and θ_2 .

5. This is a continuation of the problem discussed above. While the solution found there is formally correct based on the expansion, as with naïve perturbation theory for an anharmonic oscillator it has a problem with periodicity and a reorganized series can give much more accurate answers..
- Show on general grounds that for this problem that in the regime where system has sufficient energy to go “over the top” that $\theta(t)$ is of the form $\theta(t) = \omega t + \sum c_n \sin(n\omega t)$, where ω is in general not ω_0 .
 - Using the power counting scheme $\omega = \omega_0 + \lambda \omega_1 + \lambda^2 \omega_2 + \dots$, $c_n = \lambda^n (c_n^{(0)} + \lambda c_n^{(1)} + \lambda^2 c_n^{(2)} \dots)$ compute $\theta(t)$ up to 2nd order.
6. This is a continuation of problems 4 and 5. Consider initial conditions where the total energy E is $10V_0$. Numerically solve the equations of motion. Plot the exact solution and the 0th, 1st, 2nd order solutions for the expansions developed in 4 and 5. Briefly discuss what these plots tell you about the utility of these expansions.