

1. Problems 5.2a,b. (*Pressure gradient*) (For 5.2a, scan, photocopy, or trace the map.) For 5.2b, you'll have to work out or look up the distance across Ireland in the relevant direction.) [5+5=10 pts.]
2. Consider the function $f(x, y, z) = ax^2 + by^2 + cz^2$. [2+2+2+2+2=10 pts.]
 - (a) Find ∇f .
 - (b) Find the rate of change of f at the point $(1, 1, 1)$ in the direction of the position vector \mathbf{r} . (*Caution: \mathbf{r} is not a unit vector.*)
 - (c) Find the rate of change of f at the point $(1, 1, 1)$ in the direction of most rapid increase of f .
 - (d) The level sets $f = \text{const.}$ are ellipsoids. Find the unit normal to the ellipsoid at the point $(1, 1, 1)$.
 - (e) What is the angle between \mathbf{r} and the normal to the ellipsoid at $(1, 1, 1)$? Check that in the spherically symmetric case $a = b = c$ the angle is zero.
3. Derive the following identities [2+2=4 pts]:
 - (a) If f is a scalar field and \mathbf{v} is a vector field then

$$\nabla \cdot (f\mathbf{v}) = \nabla f \cdot \mathbf{v} + f\nabla \cdot \mathbf{v}. \quad (1)$$
 - (b) If f is a scalar field and h is a function of one variable, then

$$\nabla h(f) = h'(f)\nabla f. \quad (2)$$
4. In this problem r and \mathbf{r} are the distance and the position vector from the origin. [20 pts.]
 - (a) (i) Show using both cartesian and spherical coordinates that $\nabla r = \hat{\mathbf{r}}$. (ii) Explain why this is dimensionally balanced. (iii) Derive this equation by a geometrical discussion of the properties of the direction and magnitude of ∇r . [(2+2)+2+2=8 pts.]
 - (b) Show that $\nabla \cdot \mathbf{r} = 3$. [2 pts.]
 - (c) Show that $\nabla \cdot \hat{\mathbf{r}} = 2/r$ by the following method: write $\hat{\mathbf{r}} = r^{-1}\mathbf{r}$, and use the results of problem 3 and problem 4b. [2 pts.]
 - (d) Show that if \mathbf{m} is a constant vector, then (i) $\nabla(\mathbf{m} \cdot \mathbf{r}) = \mathbf{m}$ and (ii) $\nabla(\mathbf{m} \cdot \hat{\mathbf{r}}) = r^{-1}(\mathbf{m} - (\mathbf{m} \cdot \hat{\mathbf{r}})\hat{\mathbf{r}})$. [2+2=4 pts.]
 - (e) The magnetic field of a dipole moment \mathbf{m} is $\mathbf{B}(\mathbf{r}) = (3(\mathbf{m} \cdot \hat{\mathbf{r}})\hat{\mathbf{r}} - \mathbf{m})/r^3$. Use the above results to show that this has zero divergence (as does *any* magnetic field satisfying Maxwell's equations). (The textbook does this using the explicit cartesian components of \mathbf{B} for the case $\mathbf{m} = m\hat{\mathbf{z}}$.) [4 pts.]

(another problem follows)

5. Problems 8.2a,b,c,d (*Gravitational field of a spherically symmetric mass*) For part (c) you may just sketch the graph. No careful plot is required. [2+2+4+2=10 pts.]