

Homework Solutions, Physics 117, SPRING 2005
Home Work Problem Set # 9, (due WED 4/13/05).

James J. Griffin
301-405-6115

Page 1 of 3
Solutions by _____

Ch 11: CQ 19, 28, 40; Ex 11, 17, 19

Ch 13: CQ 1, 5 Ex 3, 7.

11 Q 19. The number of molecules is the same, by Avogadro's law that at the same temperature and pressure, EVERY LITER of ANY GAS contains the same number of molecules.

11 Q 28 ^{Let} $F = 3000 \text{ N} = P_1 A_1 = C \cdot (32 \text{ psi}) (200 \text{ cm}^2)$ describe the normal situation.

If the pressure is reduced to $P_2 = 16 \text{ psi}$ while the required Force remains the same, then the contact area must increase to A_2 as follows:

$$F_1 = P_1 A_1 = F_2 = P_2 A_2 \Rightarrow A_2 = \frac{P_1 A_1}{P_2} = \left(\frac{32}{16}\right) 200 \text{ cm}^2$$

i.e., New area $A_2 = 400 \text{ cm}^2$

11. Q 40 $T_F = 32 + \frac{180}{100} T_C$ Check that this yields $\begin{cases} T_F = 32^\circ \text{F at } T_C = 0^\circ \text{C} \\ T_F = 212^\circ \text{F at } T_C = 100^\circ \text{C} \end{cases}$

and is therefore the correct relationship between T_F & T_C

Then to get $T_F = 72^\circ \text{F}$, T_C must satisfy $72 = 32 + \frac{9}{5} T_C$

so that $T_C = (72 - 32) \cdot \frac{5}{9} = \frac{40 \cdot 5}{9} = \frac{200}{9} = \underline{22.2^\circ \text{C} = T_C}$

Homework Solutions to HW Set # 9, due WED 4/13/05.

Page 2 of 3

11. Ex 11. (a) Given Nitrogen come as N_2 , H_2 : 1 molecule has 2 atoms.
Hydrogen

Also 3 L. of H combine with 1 L. of N, & this implies that
Ammonia is $N_2 H_3$ (or $N_2 H_6$, $N_3 H_9$...), since the atom
ratio is 3:1. Then the fact that 2 L of Ammonia is produced
shows that 2 molecules of Ammonia result from every 1 molecule
of Nitrogen, which is to say from every 2 atoms of N. Thus
1 molecule of Ammonia results from each atom of N & the
chemical formula is $N_2 H_3 = NH_3$: 1 atom N + 3 atoms H
and not $N_2 H_6$, $N_3 H_9$... or any higher multiple).
Answer: 1 Nitrogen & 3 Hydrogens

11. Ex 17 $PV = nRT = \text{constant}$ when T is held constant

Then if $P \rightarrow P' = 3P$ & $V \rightarrow V' = xP$

we have $nRT = PV = P'V' = 3P \cdot xV$

$$\frac{PV}{3PV} = x = \frac{1}{3}$$

Volume is reduced by factor of 3: $V \rightarrow V' = \frac{V}{3}$
 $V_f = \frac{1}{3} L.$

11. Ex: 19 Ideal Gas Law: $P_1 V_1 = C T_1$; If temperature is kept constant while

Pressure decreases to P_2 , the new Volume is $V_2 = \frac{C T_1}{P_2}$
and the original Volume was $V_1 = \frac{C T_1}{P_1}$. Therefore

$V_2 = \frac{P_1}{P_2} V_1$. Then if $V_1 = 3 L$ at $P_1 = 100 \text{ Atm.}$ and

$P_2 = 1.25 \text{ Atm.}$, we compute $V_2 = \frac{100}{1.25} \cdot 3 L = 240 L$ at 1.25 atm

so that 240 1 L balloons can be filled.

James J. Griffin
301-405-6115

Page 3 of 3;
Solutions by

13.Q1. The gravitational potential energy is first converted to kinetic energy and then to ~~thermal~~ (heat) energy by the frictional forces which bring the avalanche to a halt.

13.Q5 Work and heat both change the internal energy of the system and can be measured in Joules. Heat operates at a microscopic level and work at a macroscopic level

13.E3. $W = Fd = 200 \text{ N} \times 4 \text{ m} = 800 \text{ J}$
 $Q = \frac{800 \text{ J} \times 1 \text{ cal}}{4.2 \text{ J}} = 190.5$

13.E7. Change in internal energy = $\Delta U = Q + W = 28 \text{ J} - 12 \text{ J}$
 $= 16 \text{ J}$
[Note Text's Answer (p A11), 164 J, is WRONG!]