Phys 410 – Homework #1

Numbered problems are from Taylor.

- 1) 1.36 (a, b and c) (9 pts).
- 2) 1.39 (9 pts).
 - a) Hint: the θ_{max} at which the range is maximized is $(\pi/4 \phi/2)$.
 - b) 3 points for proving Taylor's expression for the range, 3 points for proving my expression for θ_{max} , and 3 points for simplifying the expression for R_{max} to the result as given by Taylor.
- 3) 1.51 (numerical problem) (6 pts).
 - a) You may use any software package you prefer, including excel.
 - b) Whatever software you use, *make sure that the important constants for the problem are specified in one and only one location*, so that you can change them easily. For this problem, some important constants are, for example, g, R, the initial angle, the initial angular velocity, and the time step (Δt).
 - c) Print out and turn in the first page of your code, along with the requested plots.
- 4) 2.5 (3 pts).
 - a) You do not need to solve any equations for this question, because the solution is given in section 2.2 of the text. Instead you should describe the behavior and draw a simple sketch of the velocity as a function of time.
- 5) 2.8 (3 pts)
- 6) 2.11 (a, b, and c) (9 pts).
- 7) 2.38 (a, b, and c) (9 pts)
 - It's easiest to let (+y) point up, so that v_0 is a positive number.
 - Start with equation 2.52, except let gravity be negative. Then separate variables and integrate to find v(t).