Supplement —Phys622—Spring 1998

Prof. Ted Jacobson
Room 4117, (301)405-6020
jacobson@physics.umd.edu

Lenz vector and the Coulomb problem

Pauli found the spectrum of Hydrogen using an algebraic method. This method exploits the existence of an extra conserved quantity in the Coulomb problem: the Laplace-Runge-Lenz vector. Classically, this vector points from a focus to the perihelion along the major axis of the orbital ellipse. (See, e.g., H. Goldstein, *Classical Mechanics.*) Quantum mechanically, we define the Laplace-Runge-Lenz vector operator:

$$\mathbf{A} = \frac{1}{2m}(\mathbf{p}\times\mathbf{L} - \mathbf{L}\times\mathbf{p}) - \frac{e^2}{r}\mathbf{r}. \tag{1}$$

Since $\mathbf{A}$ commutes with the Hamiltonian H, the operator

$$\mathbf{K} = \sqrt{\frac{-m}{2H}}\,\mathbf{A} \tag{2}$$

is well-defined and Hermitian on bound states (which have negative energy). The following identities hold:

1. $\mathbf{A}$ is hermitian.
2. $[\mathbf{A}, H] = 0$.
3. $\mathbf{K}\cdot\mathbf{L} = 0 = \mathbf{L}\cdot\mathbf{K}$
4. $H = -me^4/2(\mathbf{K}^2 + \mathbf{L}^2 + \hbar^2)$
5. $[K^i, K^j] = i\hbar\epsilon^{ijk}L^k$

Defining $\mathbf{M} = (\mathbf{L}+\mathbf{K})/2$ and $\mathbf{N} = (\mathbf{L}-\mathbf{K})/2$, it follows from 3 that $\mathbf{K}^2 + \mathbf{L}^2 = 2(\mathbf{M}^2 + \mathbf{N}^2)$, and that $\mathbf{M}^2 = \mathbf{N}^2$. Also, it follows from 5, together with the angular momentum commutation relations and the fact that $\mathbf{K}$ is a vector operator, that $[M^i, N^j] = 0$ and the components of $\mathbf{M}$ and $\mathbf{N}$ each separately satisfy the angular momentum commutation relations.

Invoking the general result for the spectrum of the square of an angular momentum operator then yields the Hydrogen spectrum. The eigenvalues of $\mathbf{M}^2$ are $\hbar^2\mathcal{M}(\mathcal{M}+1)$, where $\mathcal{M}$ is an integer or half-integer, and $\mathbf{M}^2 = \mathbf{N}^2$, so 4 yields the spectrum

$$E_n = -\frac{me^4}{2\hbar^2}\frac{1}{4\mathcal{M}(\mathcal{M}+1)+1} = -\frac{me^4}{\hbar^2}\frac{1}{n^2}$$

where the principal quantum number $n = 2\mathcal{M}+1$ has been introduced. Note that although $\mathcal{M}$ can be a half-integer, n is always an integer.

The degeneracy of the level E_n is the product of the degeneracies $2\mathcal{M}+1$ for the eigenstates of $\mathbf{M}^2$ and $\mathbf{N}^2$ with eigenvalue $\hbar^2\mathcal{M}(\mathcal{M}+1)$, so the overall degeneracy is $(2\mathcal{M}+1)^2 = n^2$. Note that eigenstates of M_z and N_z are eigenstates of $L_z = M_z + N_z$, but they are *not* eigenstates of $\mathbf{L}^2$.

What about the unbound states? For these H is *positive*, so $\mathbf{K}$ as defined in (2) is *anti*-hermitian, so $\mathbf{M}$ and $\mathbf{N}$ are a combination of hermitian and anti-hermitian operators. Thus even though they satisfy the angular momentum commutation relations, we can not use the standard results for the spectrum of angular momentum operators.